






















**List of bronchodilators / inhalers to avoid or not to avoid prior to spirometry**

Bronchodilators / Inhalers <u>TO AVOID</u> before test			
Medication containing	Brand name and pictures		
<b>6 Hours</b> before test			
<b>Salbutamol</b>	Ventolin inhaler 	Salbuair inhaler 	Zenmolin tablet 
<b>Ipratropium</b>	Atrovent 	Berodual 	
<b>12 Hours</b> before test			
<b>Salmeterol</b>	Seretide Evohaler 25/50 	Seretide Evohaler 25/125 	Seretide Evohaler 25/250 
	Seretide Accuhaler 50/100 	Seretide Accuhaler 50/250 	Seretide Accuhaler 50/500 
<b>Formoerol</b>	Svmbicort 		

**List of bronchodilators / inhalers to avoid or not to avoid prior to spirometry**

<b>Bronchodilators / Inhalers <u>TO AVOID</u> before test</b>	
Medication containing <b>24 Hours</b> before test	Brand name and pictures
<b>Tiotropium</b>	Spiriva 
<b>Theophylline</b>	Neulin SR tablets 
<b><u>Umeclidinium</u></b> <b><u>Vilanterol</u></b>	Anoro® Ellipta (LAMA/LABA) 

**List of bronchodilators / inhalers to avoid or not to avoid prior to spirometry**

<b>Inhalers that <u>DO NOT</u> need to avoid prior to spirometry</b>	
<b>ASTHMA CONTROLLER INHALERS (Steroid Only) (GINA 2012 update)</b>	
<b>Brand Name DRUG and STRENGTH per Actuation</b>	<b>Brand Picture</b>
<b>BECLOMETHASONE DIPROPIONATE (HFA)</b>	
<b>Generic Beclomethasone 50mcg (HFA)</b>	
<b>Generic Beclomethasone 250mcg (HFA)</b>	
<b>BUDESONIDE</b>	
<b>Giona® Budesonide 200mcg</b>	
<b>Pulmicort® Budesonide 200mcg</b>	
<b>FLUTICASONE</b>	
<b>Flixotide® Fluticasone 50mcg</b>	
<b>Flixotide® Fluticasone 125mcg</b>	

References: Images from MIMS Online